

# Analysis of Seven Islands with Insect Fauna and Vascular Plant Flora in Gogunsan Archipelago, Korea

Jeong Ho Hwang<sup>1,2\*</sup>, Mean-Young Yim<sup>1</sup>, Seung-Lak An<sup>1</sup>, Woon-Kee Paek<sup>3</sup>, Wang-Hee Lee<sup>2</sup>

<sup>1</sup>Research and Development Division, National Science Museum, Daejeon, Korea

<sup>2</sup>Department of Smart Agriculture Systems, Chungnam National University, Daejeon, Korea

<sup>3</sup>Daegu National Science Museum, Daegu, Korea

## ABSTRACT

The analysis of seven islands in Gogunsan archipelago, Korea with insect fauna and vascular plant flora was carried out based on a field survey conducted in May, July, and September. As a result, a total of 2,817 insect individuals including 264 species and 315 taxa of vascular plant were recorded. Bangchukdo the largest island among the seven islands showed the largest number both insect species and plant taxa. The similarity analysis suggested that the nearness of each island strongly affected the insect fauna and vascular plant flora on each island. In addition, there was significant correlation between the areas of each island and the numbers of insect species (Spearman's correlation coefficient=0.857,  $P$ -value=0.014). In the future, the results of this study can be used as data related to island ecology and conservation.

**Keywords:** Archipelago, Biodiversity, Insect community, Plant community, Similarity index

## Introduction

Insects and plants are the basis of ecology and intertwined with each other through complex interactions. The relationship between insects and plant diversity has been discussed in various studies, and positive correlations have been frequently found (Gaston 1992; Procheş *et al.*, 2009; Siemann *et al.*, 1998). Plants provide a variety of benefits to insects, such as food and habitat. Therefore, an integrated investigation of these two taxa is ecologically significant and meaningful. However, although insects and plants are closely related to each other, studies

conducted simultaneous surveys of both taxa are relatively insufficient compared to individual surveys.

The ecology of island regions are geographically separated from mainland and often represents specific characteristic patterns of ecosystems, which has attracted the attention of many ecologists (Kim *et al.*, 2016). A general feature of all island systems is that they are separated from the source pool in some distance (Choi & An, 2011; Gillespie & Roderick, 2002). In this context, MacArthur and Wilson (1967) established a famous ecological theory, the theory of island biogeography. According to this theory, the biota on islands is determined by repeated inflow from mainland and extinction on islands, and biodiversity is affected by the size of the island and the distance from mainland, which is the source of species. In general, the closer to mainland and/or the larger size of an island, the greater the number of species would be found on the island (MacArthur & Wilson, 1967). Many follow-up studies on this theory have been conducted, including those that

Received October 20, 2021; Revised November 29, 2021;  
Accepted November 30, 2021

\*Corresponding author: Jeong Ho Hwang  
e-mail [chsh123@naver.com](mailto:chsh123@naver.com)  
 <https://orcid.org/0000-0001-7973-7213>



This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

applied the theory to insects or vascular plants in Korea (Jeong & Hong, 2002; Kim *et al.*, 2016; Lee *et al.*, 2008; Park *et al.*, 2013).

In Korea, approximately 4,000 inhabited and uninhabited islands are distributed in the western and southern seas, but compared to the large number of islands, the integrated database and biogeographic study of islands are insufficient (Kim *et al.*, 2016). In this context, insect fauna and vascular plant flora were simultaneously investigated and analyzed for seven islands in Gogunsan archipelago in this study. Each surveyed island was compared through similarity analysis with insects and plants, and the theory of island biogeography was applied. In addition, analyses of insect community indices and naturalized plants of each island were provided.

### Materials and Methods

#### Survey sites

The range of latitude and longitude of Gogunsan archipelago in this study was within 35.794, 126.310 to 35.862, and 126.419. Thirty-year average annual (1981–2010), maximum average annual, minimum average annual temperatures, and precipitation for this reign are 12.8°C, 17.1°C, 9.2°C, and 1,202 mm respectively (Korea Meteorological Administration, 2020). Currently, among the seven islands, Seonyudo, Jangjado, and Daejangdo are connected to mainland by bridges but Gwallido, Bangchukdo, Myeongdo, and Maldo are still separated from mainland.

The surveyed islands are as follows. Seonyu 3 district area has an area of approximately 1.17 km<sup>2</sup> in Seonyudo. Jangjado has an area of approximately 0.13 km<sup>2</sup>, which is connected to the western part of Seonyudo by the Bridge Jangjadaegyo. Daejangdo has an area of approximately 0.337 km<sup>2</sup>, which is connected to the northern part of Jangjado by the Bridge Daejangdaegyo. Gwallido has an area of approximately 1.652 km<sup>2</sup>, and is located near the western side of Jangjado and Daejangdo. Bangchukdo has an area of approximately 2.16 km<sup>2</sup> the largest island in the surveyed islands. Myeongdo has an area of approximately 0.47 km<sup>2</sup>, and is located between Bangchukdo and Maldo. Maldo is has an area of approximately 0.36 km<sup>2</sup>, and is located near the western side of Myeongdo (Fig. 1).

#### Investigation methods

Surveys were conducted for insects and vascular plants on the seven islands of Gogunsan archipelago from the 19th to the 21st of May, the 14th to the 17th of July, and the 21st, 23rd, and 24th of September 2020 (Table 1). Each of months represents spring, summer, and autumn respectively for survey the three seasons.

The survey hours for each site were approximately 2 hours during each surveyed season for both taxa. For insect surveys, sweeping and brandishing methods were used. Also, for the insects that could identified in the field were counted and recorded in the pictures. The sampled insects were identified using a stereomicroscope (Olympus SZ1145; Olympus, Tokyo, Japan). During the insect identification, various illustrated books and papers (Ahn *et al.*, 2018; Jung & Park, 2014; Lee, 2012; Park *et al.*, 2012) were referred. Entomological Society of Korean and Korean Society of Applied Entomology (1994) and National Institute of Biological Resources (2018) were used for the insect classification system. Vascular plants were photographed or cut and transferred to the laboratory for identification. For the vascular plants identification, Park (2009), Lee (2014a), Lee (2014b), Lee and Lee (2015), Cho *et al.* (2016), and Kim and Kim (2018) were used. The plant classification system was based on a checklist



Fig. 1. Survey sites in Gogunsan archipelago, Korea.

Table 1. Survey dates on each island in Gogunsan archipelago, 2020

Season	Site						
	Seonyudo	Jangjado	Daejangdo	Gwallido	Bangchukdo	Myeongdo	Maldo
Spring	2020-05-19, 20	2020-05-20	2020-05-19, 20	2020-05-21	2020-05-21	2020-05-21	2020-05-21
Summer	2020-07-17	2020-07-14	2020-07-17	2020-07-16	2020-07-16	2020-07-15	2020-07-15
Autumn	2020-09-21	2020-09-21	2020-09-21	2020-09-23	2020-09-24	2020-09-24	2020-09-23

of vascular plants in Korea (Korea National Arboretum, 2017). The sampled insects and vascular plants were stored as specimens in the Natural History Laboratory of the National Science Museum of Korea.

**Analysis methods of insect and vascular plant**

For insect community analysis, an equation transformed from the Shannon–Wiener function ( $H'$ ) derived from Margalef’s (1958) information theory was used (Pielou, 1966) for the diversity index. McNaughton’s equation (1967) was used for the dominance index, Margalef’s equation (1958) was used for the richness index, and Pielou’s equation (1975) was used for the evenness index. The similarity index was calculated using the equation developed by Sørensen (1948) and analyzed by a multi-dimensional scaling and clustering methods using IBM SPSS Statistics ver. 21 (IBM Co., Armonk, NY, USA).

Also, to analyze correlations of the distances of each island and the similarities of insect fauna and plant flora on each island, distances among the seven islands were derived from satellite images with the main surveyed areas of each island. To do this, Pearson correlations were calculated using IBM SPSS Statistics.

$$H' \text{ (Diversity index)} = -\sum \frac{n_i}{N} \ln \frac{n_i}{N} \quad (1)$$

( $n_i$ : number of individuals of the  $i^{\text{th}}$  species;  $N$ : total number of individuals)

$$DI \text{ (Dominance index)} = \frac{n_1+n_2}{N} \quad (2)$$

( $n_1$ : number of dominant species individuals,  $n_2$ : number of subdominant species individuals,  $N$ : total number of individuals)

$$RI \text{ (Richness index)} = \frac{S-1}{\ln(N)} \quad (3)$$

( $S$ : total number of species;  $N$ : total number of individuals)

$$EI \text{ (Evenness index)} = \frac{H'}{\ln S} \quad (4)$$

( $H'$ : species diversity index;  $N$ : total number of species)

$$SI \text{ (Similarity index)} = \frac{2S}{S_i+S_j} \quad (5)$$

( $S$ : number of species common to the two sites,  $S_i$ : number of species in study site  $i$ ,  $S_j$ : number of species in study site  $j$ )

To analyze naturalized vascular plants of each surveyed island, the method described by Yim and Jeon (1980) was used to calculate the urbanization indices.

In addition, according to the theory of island biogeography, the correlations among the three kinds of quantitative variables data the numbers of insect species, the numbers of plant taxa, and the areas of island were analyzed using IBM SPSS Statistics with Spearman’s correlation. The three kinds of quantitative variables data are nonparametric data and Spearman’s correlation is generally used instead of parametric Pearson’s correlation in two quantitative variables which are not normally distributed (Harris *et al.*, 2008).

The area of each island was determined based on data from the Gunsan Local Culture Encyclopedia website (GLCE, 2020). Seonyudo is originally three islands, and there are Seonyu 1, 2, and 3 districts connected by a sandbar and/or a sand dune. Seonyu 3 district area surveyed in this study was calculated after digitizing the area with QGIS based on satellite images.

**Results**

**The numbers of insect species and individuals, and vascular plant taxa**

As a result of this study, a total of 2,817 insect individuals including 264 species were collected (Supplementary Table 1). The total number of vascular plant taxa surveyed was 315 (Supplementary Table 2). Bangchukdo showed the largest numbers of insects species in any surveyed season. Vascular plants taxa also showed the largest

**Table 2.** The results of survey of insect species and plant taxa in Gogunsan archipelago, 2020

Taxa	Site							Total	
	Seonyudo	Jangjado	Daejangdo	Gwallido	Bangchukdo	Myeongdo	Maldo		
Insect	May	15 (85)	27 (77)	20 (99)	37 (74)	31 (107)	27 (167)	23 (54)	92 (663)
	July	58 (163)	28 (93)	27 (52)	58 (161)	62 (188)	58 (169)	37 (101)	142 (927)
	September	32 (72)	40 (132)	28 (150)	50 (251)	72 (205)	68 (211)	56 (206)	154 (1,227)
	Total	86 (320)	70 (302)	61 (301)	116 (486)	121 (500)	110 (547)	91 (361)	264 (2,817)
Plant	66	92	94	131	164	135	140	315	

Parenthesized numbers in ‘Insect’ row are representing the number of individuals.

**Table 3.** Dominant/subdominant insect species by the seasons and sites in Gogunsan archipelago, 2020

Season	Site									
	Seonyudo (Dominant/ subdominant species)	Jangjado (Dominant/ subdominant species)	Daejangdo (Dominant/ subdominant species)	Gwallido (Dominant/ subdominant species)	Bangchukdo (Dominant/ subdominant species)	Myeongdo (Dominant/ subdominant species)	Maldo (Dominant/ subdominant species)	Total (Dominant/ subdominant species)		
May	Megacopta punctatissima/ Aulacophora nigripennis	Megacopta punctatissima/ Syrphus ribesii, Coccinella septempunctata	Melanostoma mellinum/ Megacopta punctatissima	Megacopta punctatissima/ Aelia fieberi	Lasius japonicus/ Metasyrphus corollae	Nysius plebejus/ Pachygrontha antennata	Megacopta punctatissima/ Coccinella septempunctata	Megacopta punctatissima/ Coccinella septempunctata		
July	Nylanderia flavipes/ Pagria signata	Rivellia basilaris/ Aphaenogaster tipuna	Aphaenogaster tipuna/ Camponotus vitosus, Pagria signata	Chauliops fallax/ Monolepta pallidula	Basilepta fulvipes/ Cyphononyx fulvognathus	Bifurcanomala aulax/ Ischnura asiatica	Nysius plebejus/ Eretes griseus aulax	Basilepta fulvipes/ Bifurcanomala aulax		
September	Bothrogonia ferruginea/ Cardipennis sulcithorax	Spoladea recurvalis/ Oecanthus longicauda	Spoladea recurvalis/ Aphaenogaster tipuna	Spoladea recurvalis/ Parnara guttata	Spoladea recurvalis/ Nysius plebejus	Spoladea recurvalis/ Bothrogonia ferruginea	Locusta migratoria/ Liorhyssus hyalinus	Spoladea recurvalis/ Nysius plebejus		
Total	Megacopta punctatissima/ Aulacophora nigripennis, Bothrogonia ferruginea	Spoladea recurvalis/ Rivellia basilaris	Spoladea recurvalis/ Melanostoma mellinum	Spoladea recurvalis/ Nysius plebejus, Parnara guttata	Basilepta fulvipes/ Spoladea recurvalis	Nysius plebejus/ Pachygrontha antennata	Spoladea recurvalis/ Locusta migratoria	Spoladea recurvalis/ Megacopta punctatissima		

number in Bangchukdo among the surveyed islands (Table 2).

**Dominant and subdominant insect species by the sites and seasons**

For the total individuals of insects surveyed in this study, the dominant species was *Spoladea recurvalis*, and the subdominant species was *Megacopta punctatissima*. In September, *Spoladea recurvalis* was abundant and in May, *Megacopta punctatissima* was abundant. Jangjado, Daejangdo, and Gwallido have the same dominant species *Spoladea recurvalis*. Also, Myeongdo and Maldo have the same dominant species *Nysius plebejus* (Table 3).

**Community analysis of insect species**

As a result of insect community analysis by each island, Bangchukdo had the highest community stability, with the highest diversity, richness and evenness index, and the lowest dominance index (Fig. 2).


**The number of naturalized plant and urbanization indices of vascular plant**

As a result of the survey of naturalized plants a total of 74 taxa were found. Maldo had the highest urbanization indices among the surveyed sites (Table 4).

**Similarity analysis of insect and vascular plant**

The similarity index is regarded as a heterogeneous group when the group is less than 20% similar and a homogenous group when the group is 80% or more similar (Ahn & Park, 2012; Whittaker, 1956). Regarding the similarity analysis of insects in this study, all cases were more than 0.2, except for Daejangdo and Maldo (Fig. 3). Bangchukdo and Myeongdo had the highest similarity (0.667). The islands can be divided into two groups based on similarity indices results (Seonyudo, Jangjado, and Daejangdo group and the other islands group).

Regarding the similarity analysis of vascular plants in this study, all cases were more than 0.2 similar (Fig. 4). Bangchukdo and Maldo had the highest similarity (0.474).




**Fig. 2.** Community structure analysis of insect by the seven islands in Gogunsan archipelago, 2020. (A) Diversity index. (B) Dominance index. (C) Richness index. (D) Evenness index.

**Table 4.** The number of naturalized plant, and urbanization indices for the surveyed sites in Gogunsan archipelago, 2020

Division	Site							Total
	Seonyudo	Jangjado	Daejangdo	Gwallido	Bangchukdo	Myeongdo	Maldo	
Naturalized plant	6	20	6	28	28	16	34	74
Urbanization index	1.87	6.23	1.87	8.72	8.72	4.98	10.59	23.05

Sites	Seonyudo	Jangjado	Daejangdo	Gwallido	Bangchukdo	Myeongdo	Maldo
Seonyudo	1.000						
Jangjado	0.394	1.000					
Daejangdo	0.476	0.625	1.000				
Gwallido	0.413	0.306	0.340	1.000			
Bangchukdo	0.400	0.239	0.218	0.505	1.000		
Myeongdo	0.394	0.293	0.283	0.520	0.667	1.000	
Maldo	0.351	0.273	0.125	0.605	0.593	0.573	1.000



**Fig. 3.** Comparison of the similarity indices to insect among the sites. UPGMA, Unweighted Pair Group Method with Arithmetic Mean.

The islands can be divided into three groups based on similarity indices results (Jangjado, Daejangdo and Gwallido group, Bangchukdo, Myeongdo, and Maldo group and Seonyudo group).

To clarify correlations of the distances among the islands and the similarity indices of the two taxa, we used Pearson’s correlation analysis. As a result of analysis, the correlation coefficient between the insect similarities and the inter-island distance was  $-0.484$ , confirming the negative correlation ( $P$ -value  $0.026$ ). Also, in the case of the plant similarities and the inter-island distance, the correlation coefficient was  $-0.5$  ( $P$ -value  $0.021$ ).

**Correlation of the numbers of insect species, vascular plant taxa, and the areas of each island**


In this study, Spearman’s correlation was used to apply the theory of island biogeography. For the areas of each island and the numbers of insect species, the correlation coefficient was  $0.857$  and the  $P$ -value ( $0.014$ ) was less than significant value  $0.05$ . In the case of the numbers

of plant taxa, correlation with the areas of the island was not found. Also, the distance between the islands and mainland were not related in both the numbers of insect species and vascular plant taxa. In addition, the numbers of insect species and vascular plant taxa were analyzed to consider the relationship between insect and plant richness. The result showed the correlation coefficient of  $0.714$  and the  $P$ -value ( $0.071$ ) was larger than the significant value  $0.05$  but smaller than  $0.1$  (Table 5).

**Discussion**

The total number of surveyed insect species and individuals in this study (264 species, 2,817 individuals) was higher than that of uninhabited islands in Sinan-gun, with 122 species and 921 individuals (Cho *et al.*, 2011). These results are assumed to be because the 28 uninhabited islands in Sinan-gun are mainly smaller and the distance from mainland was larger than the islands surveyed in Gogunsan archipelago. The average area of the 28

Sites	Seonyudo	Jangjado	Daejangdo	Gwallido	Bangchukdo	Myeongdo	Maldo
Seonyudo	1.000						
Jangjado	0.316	1.000					
Daejangdo	0.400	0.441	1.000				
Gwallido	0.325	0.457	0.409	1.000			
Bangchukdo	0.339	0.414	0.380	0.468	1.000		
Myeongdo	0.348	0.335	0.341	0.398	0.468	1.000	
Maldo	0.301	0.362	0.342	0.428	0.474	0.444	1.000



**Fig. 4.** Comparison of the similarity indices to vascular plant among sites. UPGMA, Unweighted Pair Group Method with Arithmetic Mean.

**Table 5.** Spearman's correlation coefficient and *P*-value

Value	Division	
	The numbers of insect species and areas of each island	The numbers of insect species and vascular plant taxa
Correlation coefficient	0.857**	0.714*
<i>P</i> -value	0.014	0.071

\**P*<0.1; \*\**P*<0.05.

surveyed uninhabited islands in Sinan-gun is 0.048 km<sup>2</sup> which is much smaller than the average area of the seven surveyed islands in Gogunsan archipelago (0.897 km<sup>2</sup>) (Cho *et al.*, 2011).

Among the insects surveyed in this study, *Spoladea recurvalis* was the dominant species, with 341 individuals, and the subdominant species was *Megacopta punctatissima* with 147 individuals. These species were followed by *Nysius plebejus*, with 127 individuals. A survey of insects in Hallyeo Haesang National Park also confirmed that *Nysius plebejus* and *Megacopta punctatissima* were the dominant and subdominant species (Cheong *et al.*, 2005). Also, in comparison with the results of insect surveys on uninhabited islands in Sinan-gun, the subdominant species was *Spoladea recurvalis* in Lepidoptera (Cho *et al.*, 2011). The host of *Nysius plebejus* is a very wide range including Asteraceae, and the main host plant of *Mega-*

*copta punctatissima* is *Pueraria lobata* which can be commonly seen in Korea, and *Spoladea recurvalis* is a major pest with a very wide host range, so these results seems to be very general (Ahn *et al.*, 2018; Hsu & Srinivasan, 2012; Kim *et al.*, 1994).

Recently, Seonyudo, Jangjado, and Daejangdo in Gogunsan archipelago were connected to mainland by bridges. Jang and Kim (2014) conducted a vascular plants survey in Gogunsan archipelago including the same island (Jangjado) with this study before the constructions of bridges were completed and the result was that fewer vascular plants taxa were surveyed (74 taxa in Jangjado) than in this study (92 taxa in Jangjado). Also, Urbanized index for Jangjado (1.43) were lower than this study (6.23). It is assumed that the invasion of external species and moderate disturbance after the construction of the bridges caused an increase in the number of vascular plants species including naturalized plants.

By hierarchical cluster analysis, the Jangjado, Daejangdo group and the Bangchukdo, Myeongdo, and Maldo group were classified by both insects and plants suggesting high correlation between the nearness of distance and similarity indices. Also, multidimension analysis of similarity indices provided strong relation of nearness of each island intuitively because it shows similar position of islands with actual position in a satellite map in both insect and plant. The high similarity index of insects for Gwallido and Maldo may suggesting the similar environment of

two islands.

The high correlation between the numbers of insect species and the areas of each surveyed island was confirmed by Spearman's correlation coefficient. In the case of plant, the correlation between the areas and the numbers of species was not related in this study, and it is assumed that other factors such as edaphic condition, topography, sampling method, and human interference can be a greater effect in this case. Also, the distance between the island and mainland, and the numbers of species were not related in both taxa. A similar result was found in another study on vascular plant and island biogeography in Southern Sea of Jeollanamdo, Korea, reporting that this may be due to the distances between islands being too close to apply the theory of island biogeography (Kim *et al.*, 2016) which suggesting that more distances between islands should be needed to apply the island biogeography theory. For the case of the study on insect fauna in the west coast of Incheon which includes more remote islands, surveyed islands far from mainland showed less number of insect species than nearer islands (Park *et al.*, 2013).

In this study, we compared the seven islands in Gogunsan archipelago with insect fauna and vascular plant flora. Also, the correlation of the numbers of insect species, vascular plant taxa, and the areas of each island was analyzed. In the future research, it is suggested that considering more factors quantitatively such as a method of sampling, an impact of human interference, and habitat diversity. The results of this study can be used as data related to island ecology and conservation.

### Conflict of Interest

The authors declare that they have no competing interests.

### Acknowledgments

This work was supported by the National Research Foundation of Korea (grant numbers 2013M3A9A5047052, 2017M3A9A5048999).

### References

- Ahn, S.J., Kim, W.G., Kim, S.S., and Park, C.G. (2018). *The Terrestrial Heteroptera of Korea*. Seoul: Nature & Ecology.
- Ahn, S.J., and Park, C.G. (2012). Terrestrial insect fauna of the Junam wetlands area in Korea. *Korean Journal of Applied Entomology*, 51, 111-129.
- Cheong, S.W., Lee, D.J., Jeon, S.K., Kim, K.B., and Yoon, C.S. (2005). An analysis of insect community from Hallyeo Haesang National Park. *Bulletin of the Institute for Basic Science*, 17, 161-168.
- Cho, Y.H., Kim, J.H., and Park, S.H. (2016). *Grasses and Sedges in South Korea*. Seoul: GeoBook.
- Cho, Y.H., Kim, Y.J., Lim, H.M., Han, Y.G., Choi, M.J., and Nam, S.H. (2011). A faunistic study of insects of uninhabited islands in the Docho-myeon, Sinan-gun, Jeollanam-do, Korea. *Korean Journal of Environment and Ecology*, 25, 673-684.
- Choi, S.W., and An, J.S. (2011). An island network determines moth diversity on islands in Dadohaehaesang National Park, South Korea. *Insect Conservation and Diversity*, 4, 247-256.
- ESK (Entomological Society of Korea), and KSAE (Korean Society of Applied Entomology). (1994). *Check List of Insects from Korea*. Seoul: Kon-Kuk University Press.
- Gaston, K.J. (1992). Regional numbers of insect and plant species. *Functional Ecology*, 6, 243-247.
- Gillespie, R.G., and Roderick, G.K. (2002). Arthropods on islands: colonization, speciation, and conservation. *Annual Review of Entomology*, 47, 595-632.
- GLCE (Gunsan Local Culture Encyclopedia). (2020). *Geology and Topography*. Retrieved December 11, 2020 from <http://gunsan.grandculture.net/gunsan/toc?search=A1/2>.
- Harris, J.E., Boushey, C., Bruemmer, B., and Archer, S.L. (2008). Publishing nutrition research: a review of nonparametric methods, part 3. *Journal of the American Dietetic Association*, 108, 1488-1496.
- Hsu, Y.C., and Srinivasan, R. (2012). Desert horse purslane weed as an alternative host for amaranth leaf webber, *Hymenia recurvalis* in Taiwan. *Formosan Entomologist*, 32, 297-302.
- Jang, S.K., and Kim, S.W. (2014). The characteristics of flora at Gogunsan Archipelago area in Jeollabuk-do. *Journal of Korean Society for People, Plants and Environment*, 17, 37-43.
- Jeong, J.M., and Hong, K.N. (2002) Relationships between geographical conditions and distribution pattern of plant species on uninhabited islands in Korea. *The Korean Journal of Ecology*, 25, 341-348.
- Jung, B.H., and Park, H. (2014). Taxonomy of Languriinae Crotch (Coleoptera: Cucujoidea: Erotylidae) in Korea. *Korean Journal of Applied Entomology*, 53, 441-448.
- Kim, H.H., Kim, D.B., Won, H.K., Kim, C.S., and Kong, W.S. (2016). Island-biogeographical characteristics of insular flora in southern sea of Jeollanamdo, Korea. *Journal of Climate Change Research*, 7, 143-155.
- Kim, J.B., Kim, T.S., Kang, D.S., Shin, W.K., and Lee, Y.S. (1994). Investigation of pentatomid species of chrysanthemum of host plants of *Nysius Plebejus Distant* (Hemiptera: Lygaeidae) and its control. *Korean Journal of Applied Entomology*, 33, 1-5.
- Kim, T.Y., and Kim, J.S. (2018). *Woody Plants of Korean Peninsula*. Paju: Dolbegae.
- Korea Meteorological Administration (KMA). (2020). *Thirty-Year Climate*. Retrieved December 11, 2020 from <https://data.kma.go.kr/climate/average30Years/selectAverage30YearsKoreaList.do?pgmNo=188>.
- Korea National Arboretum (KNA). (2017). *Checklist of Vascular Plants in Korea*. Pocheon: Korea National Arboretum.
- Lee, C.S., and Lee, K.H. (2015). *Pteridophytes of Korea: Lycophytes & Ferns*. Seoul: GeoBook.
- Lee, H.S. (2012). *Taxonomy, phylogeny and DNA barcoding of the Korean Lauxaniidae (Diptera: Lauxanioidea)*. (Doctoral dissertation). Yonsei University, Seoul.
- Lee, S.J., Yeo, J.D., and Shin, H. (2008). Insect biogeography in the south-western Sea of Korea with comments on the insect

- fauna of Kwanmae Island. *Entomological Research*, 38, 165-173.
- Lee, T.B. (2014a). *Coloured Flora of Korea I*, 2nd ed. Seoul: Hyangmunsa.
- Lee, T.B. (2014b). *Coloured Flora of Korea II*, 2nd ed. Seoul: Hyangmunsa.
- MacArthur, R.H., and Wilson, E.O. (1967). *The Theory of Island Biogeography*. Princeton: Princeton University Press.
- Margalef, R. (1958). Information theory in ecology. *General Systems Yearbook*, 3, 36-71.
- McNaughton, S.J. (1967). Relationships among functional properties of Californian grassland. *Nature*, 216, 168-169.
- National Institute of Biological Resources (NIBR). (2018). *National Species List of Korea*. Retrieved December 11, 2020 from <http://kbr.go.kr>.
- Park, K.T., Kwon, Y.J., Park, J.K., Bae, Y.S., Bae, Y.J., Byun, B.K., et al. (2012). *Insects of Korea*. Seoul: Geobook.
- Park, S.H. (2009). *New Illustrations and Photographs of Naturalized Plants of Korea*. Seoul: Ilchokak.
- Park, S.J., Kwon, H., Park, S.K., Kim, D.S., and Park, D.S. (2013). Comparative insect faunas between Ganghwado and six other islands of west coastal in Incheon, Korea. *Journal of Asia-Pacific Biodiversity*, 6, 197-219.
- Pielou, E.C. (1966). Shannon's formula as a measure of specific diversity: its use and misuse. *The American Naturalist*, 100, 463-465.
- Pielou, E.C. (1975). *Ecological Diversity*. New York: Wiley.
- Procheş, S., Forest, F., Veldtman, R., Chown, S.L., Cowling, R.M., Johnson, S.D., et al. (2009). Dissecting the plant-insect diversity relationship in the Cape. *Molecular Phylogenetics and Evolution*, 51, 94-99.
- Siemann, E., Tilman, D., Haarstad, J., and Ritchie, M. (1998). Experimental tests of the dependence of arthropod diversity on plant diversity. *The American Naturalist*, 152, 738-750.
- Sørensen, T. (1948). *A Method of Establishing Groups of Equal Amplitude in Plant Sociology Based on Similarity of Species Content and Its Application to Analyses of the Vegetation on Danish Commons*. København: Munksgaard.
- Whittaker, R.H. (1956). Vegetation of the great smoky mountains. *Ecological Monographs*, 26, 1-80.
- Yim, Y.J., and Jeon, E.S. (1980). Distribution of naturalized plants in the Korean Peninsula. *Korean Journal of Botany*, 23, 69-83.